

# Precise spatio-temporal regulation of the anthocyanin biosynthetic pathway leads to petal spot formation in *Clarkia gracilis* (Onagraceae)

Talline R. Martins<sup>1,2</sup>, Jeremy J. Berg<sup>1</sup>, Steven Blinka<sup>1</sup>, Mark D. Rausher<sup>2</sup> and David A. Baum<sup>1</sup>

<sup>1</sup>Department of Botany, University of Wisconsin, Madison, WI, 53706, USA; <sup>2</sup>Department of Biology, Duke University, Durham, NC, 27708, USA

## Summary

Author for correspondence:

Talline R. Martins

Tel: +1 919 684 3378

Email: talline.martins@duke.edu

Received: 6 August 2012

Accepted: 19 October 2012

*New Phytologist* (2013) **197**: 958–969

doi: 10.1111/nph.12062

**Key words:** *Clarkia*, dihydroflavonol-4-reductase (*Dfr*), flower color, petal spot, pigment pattern.

- Petal spots are widespread in angiosperms and are often implicated in pollinator attraction. *Clarkia gracilis* petals each have a single red-purple spot that contrasts against a pink background. The position and presence of spots in *C. gracilis* are determined by the epistatic interaction of alleles at two as yet unidentified loci.
- We used HPLC to identify the different pigments produced in the petals, and qualitative and quantitative RT-PCR to assay for spatio-temporal patterns of expression of different anthocyanin pathway genes.
- We found that spots contain different pigments from the remainder of the petal, being composed of cyanidin/peonidin-based, instead of malvidin-based anthocyanins. Expression assays of anthocyanin pathway genes showed that the dihydroflavonol-4-reductase 2 (*Dfr2*) gene has a spot-specific expression pattern and acts as a switch for spot production. Co-segregation analyses implicated the gene products of the *P* and *I* loci as *trans*-regulators of this switch. Spot pigments appear earlier in development as a result of early expression of *Dfr2* and the flavonoid 3' hydroxylase 1 (*F3'h1*) gene. Pigments in the background appear later, as a result of later expression of *Dfr1* and the flavonoid 3'-5' hydroxylase 1 (*F3'5'h1*) genes.
- The evolution of this spot production mechanism appears to have been facilitated by duplication of the *Dfr* gene and to have required substantial reworking of the anthocyanin pathway regulatory network.

## Introduction

A major objective of evolutionary developmental biology ('Evo-Devo') is to elucidate the genetic changes that result in the evolution of novel characters. One unresolved issue is whether novel characters can arise through changes in the activity of genes that are already active in developmental programs or if they require recruitment of additional genes into those programs (Keys *et al.*, 1999). In addition, while the evolution of novel morphologies is sometimes associated with gene duplication (Xiao *et al.*, 2008; Parker *et al.*, 2009), it is unclear how frequently duplication contributes to the emergence of novel traits. Because of the apparent simplicity and ecological relevance of wing spotting patterns, the developmental control of evolutionarily novel spots in animals, particularly butterflies, has been used to address these issues (Nijhout, 1980; Carroll *et al.*, 1994; Keys *et al.*, 1999; Beldade *et al.*, 2002; Reed & Serfas, 2004). These investigations have tentatively indicated that new wing spots have arisen by the modification of existing developmental programs.

In plants, petals with large spots – discrete depositions of visible pigments that contrast with the background coloration of the flower – occur in the families Liliaceae, Orchidaceae, Asteraceae,

Papaveraceae, Fabaceae, and many others, indicating that they have evolved independently numerous times. Moreover, field studies have frequently demonstrated that spots are important in mediating interactions with pollinators (Jones, 1996; Johnson & Midgley, 1997; Van Kleunen *et al.*, 2007; Goulson *et al.*, 2009). Nevertheless, the genetic and developmental control of petal spots and, more generally, of petal color patterning has seldom been investigated. Studies in model species such as *Petunia hybrida* (petunia) and *Antirrhinum majus* (snapdragon) have clarified the genetic control of some patterns, namely vein-associated pigmentation (Schwinn *et al.*, 2006; Shang *et al.*, 2011) and variation in pigment intensity in different regions of the corolla (Jackson *et al.*, 1992; Schwinn *et al.*, 2006; Albert *et al.*, 2011). However, our general understanding of how floral pigment patterns develop and are regulated at the molecular level, specifically in relation to petals spots, remains poor. In this report, we describe experiments that begin to elucidate the biochemical, genetic and developmental basis of petal spots in the California wild flower *Clarkia gracilis*.


*Clarkia gracilis* (Onagraceae) is native to northern California, Oregon and Washington. It is the only polyploid in section *Rhodanthos*. Based on cytological analyses, Abdel-Hameed & Snow (1972) suggested that this allotetraploid derived from

hybridization between two diploid species, one closely related to *Clarkia amoena* ssp. *huntiana* and one related to *Clarkia lasenensis* and *Clarkia arcuata*. However, the parentage of *C. gracilis* has not been tested adequately by modern molecular systematic methods.

*Clarkia gracilis* is composed of four different subspecies that have somewhat overlapping ranges: *C. g.* ssp. *albicaulis*, *C. g.* ssp. *gracilis*, *C. g.* ssp. *sonomensis*, and *C. g.* ssp. *tracyi*. The subspecies differ in their floral morphologies, particularly with respect to the presence and position of petal spots. Subspecies *albicaulis* and *tracyi* have a single spot at the base of each petal, ssp. *sonomensis* has a single spot in the center of each petal, and ssp. *gracilis* lacks petal spots (Fig. 1). In all cases, mature petal spots are dark reddish-purple concentrations of pigment on a pale pink background. However, there is some variation within some subspecies: *C. g.* *sonomensis* and *C. g.* *albicaulis* can be found in either spotted or unspotted morphs.

The ecological significance of petal spots in *C. gracilis* has been studied in some detail. In mixed populations of central spotted and unspotted *C. gracilis* ssp. *sonomensis*, spotted plants have higher fitness, producing as much as 32% more seed than their unspotted counterparts in some growing seasons and sites (Jones, 1996). This difference might reflect pollinator preference, as petal spots were also shown to influence pollinator foraging behavior (Jones, 1996). In another species of *Clarkia*, *Clarkia xantiana*, the maintenance of both spotted and unspotted morphs has been attributed to pollinator-based negative frequency-dependent selection (Eckhart *et al.*, 2006).

Previous investigations have established that *C. gracilis* petal spots are under simple genetic control. Gottlieb & Ford (1988) identified two independent loci that interact epistatically to give rise to all of the morphs present in this species complex. One locus, *P*, has co-dominant alleles that determine the position of the spot within the petal: the  $P^B$  allele causes the production of basal spots, and the  $P^C$  allele causes the production of central spots. The other locus, *I*, affects the presence of basal spots: the  $I^A$  allele (or *I* in Gottlieb & Ford (1988)) suppresses basal spots, while the  $I^P$  allele (*i* in Gottlieb & Ford (1988)) allows basal spots to form. Neither allele at the *I* locus affects the formation of central spots. Gottlieb & Ford (1988) determined that *C. gracilis* ssp. *gracilis* generally has a  $P^B P^B I^A I^A$  genotype, whereas spotted plants of ssp. *sonomensis* often have the genotype  $P^C P^C I^P I^P$ . However, a variable dominance relationship was also found between  $I^A$  and  $I^P$ , suggesting that additional modifiers or alternative alleles are present in different backgrounds.



**Fig. 1** Flowers of subspecies in the *Clarkia gracilis* species complex. From left: *C. g.* ssp. *tracyi*, *C. g.* ssp. *sonomensis*, *C. g.* ssp. *gracilis*, and *C. g.* ssp. *albicaulis*.

The identity of the pigments underlying this phenotype can provide important clues as to the genes that may be involved in spot formation. Species in the genus *Clarkia* produce anthocyanin pigments, mostly derivatives of malvidin and cyanidin, although not all species produce these pigments (Dorn & Bloom, 1984; Soltis, 1986). Production of pelargonidin and derivatives is rare in the genus, being known only in *Clarkia unguiculata* (Robinson & Robinson, 1931), suggesting that the biochemical pathway branch leading to pelargonidin production is inactive in most species (Fig. 2). In *C. gracilis* flowers, the anthocyanidins malvidin, cyanidin and delphinidin (a precursor of malvidin) have been reported (Soltis, 1986).

Our objectives in this investigation were to identify genes associated with spot formation in *C. gracilis* and characterize their pattern of regulation. Specifically, we sought to answer the following questions.

- (1) What pigments are produced in *C. gracilis* flowers? Do pigments in spots differ from the rest of the petal? Do pigments in central spots differ from those in basal spots?
- (2) Is transcriptional regulation of specific anthocyanin biosynthetic genes involved in petal spot formation? If so, does any gene correspond to the spot position *P* locus?

## Materials and Methods


### Plant growth and crosses

Seeds from *Clarkia gracilis* (Piper) A. Nelson & J. F. Macbride were germinated in vermiculite at 15°C in the dark, and transferred to a 1 : 1 metromix : peatmoss mixture once seedlings had emerged. Seedlings were then transferred to 5-inch pots and were grown in a glasshouse (20–24°C) under natural light. Plants were periodically randomized to avoid possible light-induced changes in flower color.

Several individuals from various populations were grown, and only those from monomorphic (with respect to spots) populations were used as the parents for crosses. Two independent crosses were carried out to separate the *P* and *I* loci. For the ‘*P*’ cross, a central-spotted *C. g.* *sonomensis* was crossed to a basal-spotted *C. g.* *albicaulis*. For the ‘*I*’ cross, an unspotted *C. g.* *gracilis* was crossed to a basal-spotted *C. g.* *albicaulis*. Voucher information is available in Supporting Information Table S1. Pollen recipient flowers were emasculated several days before full stigma receptivity to avoid self-pollination. Stigmas were pollinated by touching them with a dehiscing anther from the designated sire. F1 individuals were selfed to obtain F2 progeny.

### Pigment extraction and high-performance liquid chromatography (HPLC) analyses

Anthocyanins were extracted from F2 individuals derived from the ‘*P*’ cross mentioned in the previous section as well as from *C. gracilis* individuals derived from several natural populations. Seven F2 individuals were examined: two central-spotted, two basal-spotted, and three double-spotted. Five to ten F1 individuals (all double-spotted) were pooled into one sample. Individuals



**Fig. 2** Simplified schematic representation of the anthocyanin biosynthetic pathway. Abbreviations for enzymes: CHS, chalcone synthase; CHI, chalcone isomerase; F3H, flavanone 3 hydroxylase; F3'H, flavonoid 3' hydroxylase; F3'5'H, flavonoid 3'-5' hydroxylase; DFR, dihydroflavonol-4-reductase; ANS, anthocyanidin synthase; FLS, flavonol synthase; UF3GT, UDP glucose:flavonoid 3-O-glucosyltransferase. Dihydroflavonols: DHK, dihydrokaempferol; DHQ, dihydroquercetin; DHM, dihydromyricetin.

from natural populations included three *C. g. ssp. albicaulis* and six *C. g. ssp. sonomensis* plants (three spotted and three unspotted; see Table S1 for voucher information). Each sample contained the top, center, or base of four to five adult flowers per plant (Fig. S1).

Anthocyanidins were extracted as in Harborne (1984). Pigments were separated using HPLC and identified using the retention times of commercially available standards (delphinidin, pelargonidin, cyanidin, peonidin, malvidin, and petunidin; Polyphenols Laboratory, Sandnes, Norway) that were run using the same conditions as the unknown samples (previously described in Smith & Rausher, 2011).

#### DNA extraction, gene amplification and sequencing

Genomic DNA from *C. gracilis* ssp. for gene isolations and phylogenetic analyses was extracted from fresh plant tissue using the Cetyltrimethylammonium bromide (CTAB) method (Doyle & Doyle, 1987). Genomic DNA from *C. amoena* ssp. *huntiana*, *C. lassenensis*, *C. arcuata*, *Clarkia franciscana*, *Clarkia rubicunda*, and *Clarkia breweri* was kindly provided by Kenneth Sytsma (University of Wisconsin-Madison; voucher information in Table S1). We identified the two homeologs of *C. gracilis* genes (arbitrarily called *A* and *B*) by including sequences from

other species in section *Rhodanthos* in phylogenetic analyses. This was facilitated by *C. amoena* and *C. lassenensis* (the putative parents of *C. gracilis*) not being sister taxa, allowing the homeologs present in *C. gracilis* to be identified more easily.

We used sequences available in GenBank to design degenerate primers to isolate partial sequences corresponding to the anthocyanin biosynthetic genes (flavanone 3 hydroxylase (*F3h*), flavonoid 3' hydroxylase (*F3'h*), flavonoid 3'-5' hydroxylase (*F3'5'h*), dihydroflavonol-4-reductase (*Dfr*) and anthocyanidin synthase (*Ans*)). Primers and conditions used for each gene/species are available upon request. PCR bands corresponding to predicted amplicon sizes were transformed into *Escherichia coli* cells. At least 10 colonies per ligation were sequenced, and additional clones were sequenced as needed. For sequencing reactions, the BigDye Terminator v. 3.1 system was used following the manufacturer's protocol (Applied Biosystems, Foster City, CA, USA) and electrophoresis was performed at the University of Wisconsin-Madison Biotechnology Center.

#### Sequence and phylogenetic analyses

SEQUENCHER v. 4.7 (Gene Codes Corp. Ann Arbor, MI, USA) was used to edit, assemble and align sequence fragments with

further manual alignment in MACCLADE v. 4.05 (Maddison & Maddison, 2000). To confirm gene identity, sequences were submitted to the Basic Local Alignment Search Tool at the National Center for Biotechnology Information. All phylogenetic analyses were performed in PAUP\* v. 4.0b10 using maximum likelihood (ML; Swofford, 2002). Potential PCR recombinants were excluded from the analyses. For all data sets, the generalized time reversible (GTR) model with estimated base frequencies was assumed. Support values were obtained by performing 1000 replicates of ML bootstrapping. Only coding sequences were used in analyses (alignments of included sites are provided in Notes S1). Sequences isolated in this study can be found under GenBank accession numbers KC019211–KC019249.

### mRNA expression analyses

Total RNA was extracted from petals using the RNeasy Plant Mini kit (Qiagen, Hilden, Germany), followed by a DNase step (MBI Fermentas, St Leon-Rot, Germany) and cDNA synthesis using the ImpromII Reverse Transcription System (Promega, Madison, WI, USA), or Superscript III First Strand Synthesis (Invitrogen, San Diego, CA, USA), all according to the respective manufacturer's protocol.

For qualitative gene expression experiments, F2 plants of three different phenotypes (central-, basal-, and double-spotted) from the *P* locus cross and of two different phenotypes (basal-spotted and unspotted) from the *I* locus cross were used. Each RNA sample consisted of either the top, base, or center of three to four flower buds from each plant and three plants for each phenotype. Buds were collected at a stage of development when spot pigments were first appearing (or in equivalently sized petals of unspotted plants), but before background color was visible (Fig. S2).

For quantitative gene expression experiments, plants of three different phenotypes (central-, basal-, and unspotted), derived from the *P* and *I* crosses, were used. Buds representing four different stages of development were collected, with three replicates of each stage. 'Stage' was based on the emergence and quantity of pigments, because petal length proved to be an unreliable marker for age, as flower size and maturation times varied greatly, within and between individuals. Change in floral pigmentation has been used previously in *Clarkia* spp. to determine floral developmental stage (Pichersky *et al.*, 1994). In *C. gracilis*, color develops in the following order: no color, central spot appears (if present), basal spot appears (if present), and background color appears (Fig. S3).

Quantitative assays of gene expression were carried out using Clontech's (Clontech, Mountain View, CA, USA) SYBR Advantage qPCR premix and run in a Stratagene (Stratagene, La Jolla, CA, USA) Mx3000P instrument. Dissociation curves and sequences of qPCR products were obtained to verify amplification of only the desired product. Absolute quantification of mRNA transcript levels was carried out using a standard curve, followed by normalization to *Actin*. qPCR runs were performed in duplicate. Primers used for RT-PCR and qPCR (Table S2) flanked introns, with the exception of *F3h* and *Ans*. PCR products were sequenced to confirm that the desired copy was being amplified.

### Genotyping F2 individuals from P and I crosses at the *Dfr2-A* and *Dfr2-B* loci

Thirty-two and 31 F2 plants from the *P* cross were genotyped for *Dfr2-A* and *Dfr2-B*, respectively. Homeolog-specific primers were designed to amplify a portion of exon 2 and intron 2 (*Dfr2-A*), and a portion of intron 2 (*Dfr2-B*). Fifty plants from the F2 generation from the *I* cross were genotyped at *Dfr2-A* and *Dfr2-B* by amplifying intron 3 and partial sequences of exons 3 and 4. The same *Dfr2* primers used for qPCR analyses were used. All alleles were identified using fragment length polymorphisms, with the exception of alleles of *Dfr2-B* for the *P* cross, which were genotyped by the presence of a single nucleotide polymorphism (SNP) using restriction enzyme digestion (Tables S3, S4). Fragment length runs were completed in an ABI 3730XL DNA Analyzer at the Duke Genome Sequencing & Analysis Core Resource, visualized using the GENEMAPPER software (Applied Biosystems, Foster City, CA, USA) and scored by eye. Fisher's exact test was used to test for possible co-segregation between spot presence and *Dfr2-A/B* alleles.

### Results

#### Crosses confirm the two-locus model and provide segregating families for further analysis

Although the genetics of petal spot inheritance had already been determined, we established populations of individuals with known genotypes to facilitate genetic and molecular analyses and to confirm previous genetic results. We developed two separate segregating F2 populations for the two independent loci, thus removing the epistatic interaction that could mask the genotype at the *P* locus.

Segregation patterns (Table 1) confirmed those previously reported (Gottlieb & Ford, 1988; Jones, 1996). In the *P* cross (see Materials and Methods), 100% of F1 progeny were double-spotted. Segregation in the F2 population did not deviate significantly from the expected 1 : 2 : 1 ratio of basal-spotted:double-spotted:center-spotted plants ( $\chi^2$  test,  $P=0.74$ ) consistent with a model of a single locus with co-dominant alleles. According to this model,  $P^B P^B$  plants have basal spots,  $P^C P^C$  plants have central spots, and  $P^B P^C$  plants have both a central and a basal spot.

**Table 1** Segregation for presence vs absence and position of petal spot in two separate crosses, in F1 and F2 generations of *Clarkia gracilis* plants

|                |     | Spot position |         |       |        |
|----------------|-----|---------------|---------|-------|--------|
|                |     | <i>n</i>      | Central | Basal | Double |
| <i>P</i> cross |     |               |         |       |        |
| F1             | 23  |               | –       | 23    | –      |
| F2             | 82  | 18            | 23      | 41    | –      |
| <i>I</i> cross |     |               |         |       |        |
| F1             | 7   | –             | 7       | –     | –      |
| F2             | 101 | –             | 77      | –     | 24     |

*P* cross refers to central-spotted *C. g. sonomensis* × basal-spotted *C. g. albicaulis* cross, while *I* cross refers to unspotted *C. g. gracilis* × basal-spotted *C. g. albicaulis* cross.

In the *I* cross (see Materials and Methods), all F1 individuals had basal spots, suggesting that the suppressing  $I^A$  allele was recessive to the spot-permitting allele  $I^P$ . In the F2 population, we saw variation in spot intensity, from a strongly defined spot to an ill-defined 'smudge', to unspotted petals. Because it was difficult to distinguish spots from smudges, plants showing any color at the base were coded as 'basal-spotted' for statistical analyses. As shown in Table 1, using this scoring, a 3 : 1 pattern of spotted: unspotted plants could not be rejected ( $\chi^2$  test,  $P=0.77$ ), suggesting that the  $I^P$  allele isolated is generally (but variably) dominant to the  $I^A$  allele. Genotypes for *P* and *I* and expected phenotypes under this genetic model are presented in Fig. S4.

It should be noted that, while in our crosses the  $I^P$  allele appeared to be dominant, previous work has shown that alleles at the *I* locus can switch from dominant to recessive in different backgrounds, possibly due to other modifiers (Gottlieb & Ford, 1988). Further genetic studies are necessary to fully understand the phenotypic expression of this allele in different *C. gracilis* populations and subspecies.

#### HPLC analyses show that spots contain different pigments from the rest of the petal

While the pigment composition of entire *C. gracilis* petals had been previously determined through chromatography (Soltis, 1986), it was unclear whether the spots contained the same pigments as the background color. Using HPLC, we analyzed anthocyanidins present in mature petals from several *C. gracilis* individuals, spanning all spot phenotypes (see Materials and Methods). In all cases, irrespective of subspecies or spot phenotype, the distribution of anthocyanidins in the petal followed the same pattern. Petals of *C. gracilis* were found to include the anthocyanidins malvidin, cyanidin, and peonidin (Fig. 3, Table 2). These results contrast with a published report where delphinidin was found instead of peonidin (Soltis, 1986). The discrepancy between this study and Soltis (1986) may be a result of the different methodologies used (HPLC vs thin-layer chromatography).

Cyanidin and peonidin were present in all petal sections that included spots and were never detected in petal sections that lacked spots (Fig. 3, Table 2). Likewise, petal sections that lacked spots, only having background color, contained only malvidin. The central regions of central-spotted flowers sometimes

contained small amounts of malvidin, consistent with the presence of some background color in the central sector. In contrast, the basal regions of basal-spotted flowers contained very little or no malvidin, consistent with the almost complete lack of background color near the petal base (Fig. S1).

Given these findings, we conclude that, regardless of spot position, anthocyanins in *C. gracilis* petal spots are derived from cyanidin and peonidin, two similar anthocyanidins that differ in that peonidin has a methyl substitution on the B-ring. In contrast, the anthocyanins present in the background of the petal appear to be derived from malvidin, an anthocyanidin that has two methyl substitutions on the B-ring and derives from a different branch of the anthocyanin pathway than cyanidin and peonidin (Fig. 2).

#### *Clarkia gracilis* carries multiple copies of anthocyanin biosynthetic genes


In the anthocyanin pathway the branch-point between cyanidin/peonidin and malvidin occurs after the F3H-mediated synthesis of dihydrokaempferol (DHK) (Fig. 2). This indicates the regulation or action of *F3'h* and/or *F3'5'h* as possible determinants of spot formation. Another possible candidate for the synthesis of different pigments is *Dfr*. Numerous studies show that single or very few amino acid changes in DFR can lead to changes in substrate specificity that redirect flux down a different branch of the pathway (Johnson *et al.*, 2001; Fischer *et al.*, 2003; Shimada *et al.*, 2005; DesMarais & Rausher, 2008). Consequently, we focused on documenting whether differences in the patterns of regulation of *F3'h*, *F3'5'h* and *Dfr* are associated with the different petal spot phenotypes of *C. gracilis*. We also examined *F3h* and *Ans*, which are involved in earlier and later steps of pigment biosynthesis, respectively, to serve as controls and provide a more comprehensive understanding of patterns of expression of anthocyanin biosynthetic genes.

We used specific and degenerate primers to amplify partial sequences and cloned numerous PCR products in an attempt to identify all homologous gene copies in *C. gracilis* and, when possible, in related diploid species. We analyzed sequences for the presence of stop codons to assess gene functionality and used phylogenetic analyses to confirm gene identity (Fig. S5). However, it should be noted that we obtained only partial sequences for most genes from most species and that, due to the lack of a sequenced genome, gene copy number could not be determined unequivocally.

**Table 2** Pigment composition in *Clarkia gracilis* petals with different spot phenotypes

|               |        | Phenotype               |                         |                         |                         |
|---------------|--------|-------------------------|-------------------------|-------------------------|-------------------------|
|               |        | Basal spot              | Central spot            | Central and basal spots | No spot                 |
| Petal section | Top    | 0.171 ( $\pm$ 0.050), 5 | 0.132 ( $\pm$ 0.026), 3 | 0.205 ( $\pm$ 0.102), 3 | 0.137 ( $\pm$ 0.033), 2 |
|               | Center | 0.227 ( $\pm$ 0.078), 2 | 3.524 ( $\pm$ 1.599), 5 | 2.717 ( $\pm$ 0.179), 3 | 0.143 ( $\pm$ 0.026), 2 |
|               | Base   | 4.55 ( $\pm$ 1.020), 5  | 0.494*, 1               | 5.316 ( $\pm$ 1.091), 3 | 0.355 ( $\pm$ 0.115), 2 |

Numbers refer to the ratio of cyanidin + peonidin : malvidin (mean  $\pm$  standard deviation), followed by number of individuals sampled. The amount of each pigment was inferred from the average peak height between 10.4 and 10.5 min for cyanidin, between 11.1 and 11.2 min for malvidin, and between 12.3 and 12.4 min for peonidin. \*All but one central-spotted individual had spots that extended close to the petal base (on the abaxial but not the adaxial surface), impeding dissection of spot-free tissue. Therefore, only one individual was used.




**Fig. 3** Anthocyanidins found in *Clarkia gracilis* flowers. HPLC traces show pigments in different sections of petals. Each panel (a–d) shows traces for sections of the petal (top, center, and base, in that order) for one of the four phenotypic classes: basal-spotted (a); central-spotted (b); double-spotted (c), and unspotted (d). Peaks corresponding to malvidin, cyanidin, and peonidin (indicated x, y, and z, respectively) were identified by comparison to known standards.

Two copies of *F3h*, *F3'h*, and *F3'5'h* and three copies of *Dfr* were found in diploid species in Sect. *Rhodanthos*, while a single copy of *Ans* was found. For *F3'h* and *F3'5'h*, only one copy appeared to be functional in the diploids: one copy of each gene had a premature stop codon or a frameshift mutation. For *F3h*, one copy from *C. amoena* and one from *C. arcuata* appeared to be divergent (as evidenced by longer branches; see Fig. S5). There were no indications from the sequences to suggest that any *Dfr* genes lack function (Fig. S5). We focused our efforts on isolating the putative functional or nondivergent copies of these genes from *C. gracilis*, and recovered one copy (two homeologs) corresponding to the putative functional *F3h* and *F3'h*, and all three copies (and two homeologs for each copy) of *Dfr*. We isolated both homeologs of the functional copy of *F3'5'h*, and one homeolog of the nonfunctional *F3'5'h* gene, which was confirmed to have an early stop codon. We isolated a single sequence corresponding to *Ans*, but it is unclear whether only one homeolog exists, or whether both homeologs are present but do not differ in sequence within the short fragment obtained.

#### Spatial expression of *Dfr2* shows a tight correlation to spot location in *C. gracilis* petals

If spot formation depends on location-specific expression of any of the anthocyanin biosynthetic genes, it follows that such gene (s) would be expressed mainly in areas where spots are formed. To examine spatial expression patterns, we performed RT-PCR on RNA extracted from three different petal sections (top, middle and base) from young buds, from plants with central- ( $P^C P^C$ ), basal- ( $P^B P^B$ ), or double-spotted ( $P^B P^C$ ) flowers. Copy-specific primers were used for all of the putatively functional genes, with homeolog-specific primers being used in *C. gracilis* when that proved possible.

As shown in Fig. 4, the gene encoding the most upstream enzyme examined in the anthocyanin pathway, *F3h*, and one of the two copies of the gene encoding the next enzyme, *F3'h1-A*, showed consistent, robust expression in all sections of all petals. *F3'h1-B* transcript was not found in floral tissue, despite multiple attempts using several primers, suggesting that it is either not



**Fig. 4** Spatial expression of anthocyanin biosynthetic genes in *Clarkia gracilis* petals. (a) Expression patterns for three individual plants for each of the *P* locus phenotypes. The same nine individuals were used for all genes, and they are placed in the same position for each panel. Ovals represent petals, and filled circles represent spot locations. T, C, and B refer to top, central, and basal sections of the petals, respectively. *F3h*, flavanone 3 hydroxylase; *F3'h*, flavonoid 3' hydroxylase; *F3'5'h*, flavonoid 3'-5' hydroxylase; *Dfr*, dihydroflavonol-4-reductase; *Ans*, anthocyanidin synthase. (b) Expression patterns for *Dfr2* in three spotted and three unspotted plants from an F2 population segregating for *I*.

expressed or limited to other tissues. Genes encoding enzymes further down the pathway, such as *F3'5'h1*, *Dfr1*, and *Ans* (hereafter, 'downstream genes'), showed variable expression that did not correlate with genotype or spot location. This variation probably reflects slight differences in the age of petals. For instance, the most consistent pattern among the downstream genes was that their levels tended to be lower in central-spotted petals than in either basal-spotted or double-spotted petals. This


was probably a consequence of the age of the floral buds. Central spots appeared earlier in development than basal spots, which meant that central-spotted flowers were harvested at a slightly earlier stage of development than the other genotypes (Fig. S3). This is significant because it suggests that, at this early stage of development, downstream genes are not fully and consistently activated.

The two homeologs of one gene, *Dfr2*, had a pattern that was strikingly distinct from all the other genes surveyed, a pattern that was correlated with the presence of the spot (Fig. 4a). *Dfr2* appeared to be expressed at high levels in sections of the petal that contained a spot, but not in unspotted sections. In central-spotted individuals *Dfr2* was expressed at high levels only in the center of the petal, but was expressed only at the base of basal-spotted petals. Similarly, double-spotted individuals showed high expression of *Dfr2* in both central and basal sections of the petal. The low levels of expression that were sometimes seen in sections adjacent to the spot can be attributed to imprecise sectioning of the buds, as they are only a few millimeters in length when dissected. The sole exception to this pattern is that one central-spotted individual showed a brighter band for *Dfr2* in the top as opposed to the central section of the petal (Fig. 4a). This could be partly attributable to initial RNA quality, as the top section for that petal yielded a brighter band than the central sections for the other genes surveyed, including *Actin*. The spot-specific pattern of expression of *Dfr2* in petals (and the lack of *Dfr2* expression in unspotted petals) was confirmed in nine other *C. gracilis* plants grown from seed derived from wild populations (three unspotted and two central-spotted *C. g. sonomensis* plants, three basal-spotted *C. g. albicaulis* plants, and one basal-spotted *C. g. tracyi* plant; Fig. S6). Therefore, we conclude that *Dfr2* (and only this gene) shows a spot-specific expression pattern.

The role of *Dfr2* in spot formation was further evaluated by examining the expression of this gene in individuals segregating for the *I* locus. In basal-spotted plants ( $I^P I^P$  or  $I^P I^A$ ), *Dfr2* was expressed at high levels in the basal section, whereas unspotted siblings ( $I^A I^A$ ) showed very low or no *Dfr2* expression (Fig. 4b). This corroborates the pattern seen in F2s from the *P* locus cross, suggesting that *Dfr2* is necessary for spot production. It also indicates that suppression of the basal spot by the *I* locus is genetically upstream of *Dfr2*.

#### *Dfr2* deviates from other downstream genes in showing expression early in petal development

The variable expression of genes encoding downstream enzymes suggested the possibility that temporal regulation plays a role in petal spot development. This seems plausible as petal spots, especially central spots, are visible several days before the appearance of background color (Fig. S3), implying that genes required for spot development might be expressed earlier than genes that are only required for the production of background color. To investigate this possibility, we used qPCR to obtain an expression time-course for *Dfr* and for the functional copies of *F3'h* and *F3'5'h*, which are situated at the branch-point between cyanidin/peonidin and malvidin biosynthesis in developing buds. We assayed



**Fig. 5** Quantitative analysis of gene expression of anthocyanin biosynthetic genes through *Clarkia gracilis* petal development. Photographs above the graphs show a representative petal for each of the four developmental stages defined for the purpose of this experiment. From L to R: basal, central, and unspotted. All buds are shown in the upright position (apex at the top; base at the bottom). From left to right, basal-spotted plants: spot first appearing, spot well defined, background beginning to show, and mature petal; central-spotted plants: no color, spot first appearing, spot well formed, and background color beginning to appear; unspotted plants: young colorless, older colorless, background first appearing, and mature petal. *y*-axes show expression of genes normalized to *Actin*. *F3'h*, flavonoid 3' hydroxylase; *F3'5'h*, flavonoid 3'-5' hydroxylase; *Dfr*, dihydroflavonol-4-reductase. Columns represent averages across three replicates (except the last time-point for central-spotted; *n* = 2), and error bars indicate standard error.

all three copies of *Dfr* and both homeologs for all genes except the *B* copy of *F3'h1*, which did not appear to be expressed in petals.

As shown in Fig. 5, two genes, *F3'h1-A* and *Dfr2*, were expressed early in development, having high expression levels at the first time-points (as the spot appeared) and declining expression in more mature buds (which were developing background color). Although *F3'h1-A* showed a similar pattern in all three phenotypes, *Dfr2* expression was reduced or absent in flowers that lacked spots, consistent with the results obtained from non-quantitative PCR (Fig. 4). In contrast to *F3'h1-A* and *Dfr2*, *F3'5'h1* and *Dfr1* were expressed later in petal development, having low expression levels at early time-points, but higher expression later in development.

Finally, one gene, *Dfr3*, showed no clear pattern, being expressed at very low levels in all tissues except for a peak in late, central-spotted flowers. The latter could be an experimental artifact (note error bars, Fig. 5). The success in amplifying *Dfr3* in RT-PCR experiments confirmed that this gene was expressed, but the level of expression appeared to be substantially lower than

that of the other *Dfr* gene copies, indicating that this gene may lack a function in pigment production in the petal.

#### Co-segregation analysis suggests that the *P* and *I* loci correspond to *trans*-regulators of *Dfr2*

Qualitative and quantitative gene expression data indicate that cyanidin/peonidin-based petal spots in *C. gracilis* correlate with the early and localized expression of *Dfr2*, coupled with the early expression of *F3'h1-A*. Under the assumption that localized, early expression of *Dfr2* is the proximate cause of petal spot development, the *P* and *I* loci must act by modulating the expression of *Dfr2*, and probably correspond to upstream regulators of *Dfr2*. However, it is possible that either *P* or *I* could correspond to *Dfr2* itself if it had undergone a *cis*-regulatory change that resulted in altered expression.

To verify whether *Dfr2* alleles co-segregate with spot location and/or presence, we genotyped several F2 individuals from both the *P* and *I* crosses, for both homeologs of *Dfr2* (Table 3). The genotypes at neither of the *Dfr2* homeologs co-segregated

perfectly with spot location in the *P* cross, suggesting that the *P* locus does not encode *Dfr2*, but rather is a *trans*-regulator of *Dfr2*. However, the *P*-value for co-segregation between *Dfr2-B* and *P* approached significance ( $P=0.0843$ ), suggesting that perhaps there is some genetic linkage between *P* and *Dfr2-B*. Larger numbers of offspring would be necessary to confirm this hypothesis. Similarly, neither of the *Dfr2* homeologs co-segregated with the absence/presence of basal spots in F2s from the *I* locus cross, suggesting that *I* is also a *trans*-regulator of *Dfr2*.

## Discussion

### A model for the development of pigmentation patterns in *C. gracilis* petals

Based on the results presented here, we propose a model for petal pigment development in *C. gracilis* that can explain the production of cyanidin/peonidin-based anthocyanins in spots and malvidin-based anthocyanins in the background (Fig. 6). This model invokes transcriptional regulation of anthocyanin biosynthetic genes as a function of time and space. Although post-translational regulation may occur, it is not necessary to explain the observed data. For this model, we assume that *Dfr3* is expressed at such low levels that it plays no significant role in petal pigmentation. Likewise, we will ignore the putatively nonfunctional copies of *F3'h* and *F3'5'h*.

According to the model (Fig. 6), most of the core enzyme-coding genes of the anthocyanin pathway (chalcone synthase (*Chs*), chalcone isomerase (*Chi*), *F3h*, *F3'h*, *Ans*, and UDP glucose:flavonoid 3-O-glucosyltransferase (*Ufgt*)) are activated early in development and expressed throughout petal tissue. In addition, *Dfr2* is activated early but is expressed only in the area destined to become a pigmented spot, implying that it is under different regulatory control. By contrast, neither *F3'5'h1* nor *Dfr1* is expressed at this time. Because *Dfr* is required for pigment production, the spatial restriction of *Dfr2* explains why pigment is produced only as a spot. Further, because *F3'5'h1* is not activated early, only anthocyanins derived from dihydroquercetin (DHQ) (i.e. cyanidin and peonidin) are produced in the spot domain.

At a later stage of development, the core enzymes are still active. This may represent continuous expression of these genes, or a second wave of expression. At this point, *F3'h1-A* and *Dfr1* expression has ceased. Activation of both *F3'5'h1* and *Dfr1* throughout the developing petal means that the complete set of enzymes is present, causing the background pigment to be deposited. The expression of *F3'5'h1* causes this pigment to be derived via dihydromyricetin (DHM) rather than DHK or DHQ, explaining the production of malvidin in the background. Thus, the difference in pigment type between background and spot, as well as the spatial restriction of the spot, are explained by the different temporal expression of *F3'h1-A/Dfr2* vs *F3'5'h1/Dfr1*.

### Regulation of enzyme-coding genes

In most species that have been examined, the core enzyme genes of the anthocyanin pathway (those depicted in Fig. 2) are coordinately regulated by common sets of transcription factors (Cone *et al.*, 1986; Paz-Ares *et al.*, 1986, 1987; Ludwig *et al.*, 1989; Goodrich *et al.*, 1992; de Vetten *et al.*, 1997; Elomaa *et al.*, 1998; Quattrocchio *et al.*, 1998, 1999; Walker, 1999; Borevitz *et al.*, 2000; Spelt *et al.*, 2000, 2002; Carey *et al.*, 2004; Schwinn *et al.*, 2006; Gonzalez *et al.*, 2008; Lin-Wang *et al.*, 2010; Niu *et al.*, 2010; Albert *et al.*, 2011). In some taxa, such as *Ipomoea* and maize (*Zea mays*), all core genes are expressed at similar times in development and are activated by a single set of transcription factors (Cone *et al.*, 1986; Paz-Ares *et al.*, 1987; Ludwig *et al.*, 1989; Morita *et al.*, 2006). In other species, such as *Petunia hybrida* and *Antirrhinum majus*, genes are activated in two blocks, with those coding for upstream enzymes activated first, and those coding for downstream enzymes activated later, perhaps by a different set of transcription factors (Martin *et al.*, 1991; Goodrich *et al.*, 1992; Jackson *et al.*, 1992; Quattrocchio *et al.*, 1998; Spelt *et al.*, 2000; Schwinn *et al.*, 2006).

The pattern of regulation exhibited by *C. gracilis* differs from these canonical patterns. Although we have not demonstrated it directly, the production of anthocyanins in the petal implies that upstream and downstream genes, represented by *F3h* and *Ans*, are expressed from the time anthocyanins are first produced in spots to the time they are last produced in the remainder of the

**Table 3** Analysis of co-segregation of alleles at the dihydroflavonol-4-reductase 2 (*DFR2*)-*A* and *DFR2*-*B* loci with phenotypes obtained from the *I* and *P* crosses

| Homeolog      | Genotype | <i>I</i> Cross |          |          | <i>P</i> Cross |          |          |          |
|---------------|----------|----------------|----------|----------|----------------|----------|----------|----------|
|               |          | Phenotype      |          |          | Phenotype      |          |          |          |
|               |          | <i>S</i>       | <i>U</i> | <i>P</i> | <i>B</i>       | <i>D</i> | <i>C</i> | <i>P</i> |
| <i>Dfr2-A</i> | A1A1     | 11             | 2        |          | 3              | 5        | 1        |          |
|               | A1A2     | 19             | 5        | 0.591    | 5              | 8        | 5        | 0.9314   |
|               | A2A2     | 12             | 1        |          | 2              | 2        | 1        |          |
| <i>Dfr2-B</i> | B1B1     | 10             | 4        |          | 3              | 1        | 0        |          |
|               | B1B2     | 18             | 3        | 0.2365   | 0              | 6        | 2        | 0.0843   |
|               | B2B2     | 14             | 1        |          | 6              | 8        | 5        |          |

Different alleles of *Dfr2-A/B* were identified using single nucleotide polymorphisms (SNPs) and indels in intron 2 or 3. *S*, spotted; *U*, unspotted; *B*, basal-spotted; *D*, double-spotted; *C*, central spotted. *P*-values are from Fisher's exact test.

petal—although the possibility of the existence of sub-blocks within these genes cannot be ruled out. However, *Dfr*, *F3'h*, and *F3'5'h* exhibit more restricted temporal distributions, with *Dfr2* and *F3'h1-A* being expressed only in the early stages of development and *Dfr1* and *F3'5'h1* being expressed only later in development. This pattern suggests that factors regulating these genes differ from those regulating the remaining core enzymes, and that the factors regulating *Dfr2* and *F3'h1-A* differ from those regulating *Dfr1* and *F3'5'h1*. Moreover, two patterns indicate that *Dfr2* is regulated differently from all other enzyme-coding genes. First, only *Dfr2* exhibits a spatially restricted expression domain, confined to the regions that develop spots. Secondly, variation at the *P* and *I* loci affects only expression of *Dfr2*. Because neither of these loci co-segregates with *Dfr2*, they probably correspond to transcription factors or cofactors that are unique to this gene. These patterns imply that the production of spots has required a substantial reworking of the anthocyanin regulatory network. We next discuss the evolutionary implications of this reworking.


#### Possible substrate specificity in different copies of DFR

DFR2 and DFR1 appear to only encounter one substrate, DHQ or DHM, respectively, and, while the model we propose for formation of cyanidin/peonidin-based spots does not require evolution of substrate specificity among the different copies of DFR, it is possible that it has occurred. In *C. gracilis*, the two different copies differ in numerous sites, including in the active site, where they differ by nine amino acid substitutions. One such substitution, at residue 133 (Fig. S7), has been implicated in substrate specificity in other species (Johnson *et al.*, 2001; Shimada *et al.*, 2005; DesMarais & Rausher, 2008). Possible evolution of substrate specificity could be explored by biochemical characterization of these enzymes combined with more fine-scale temporal expression studies.

#### Evolutionary implications

Although the progenitors of *C. gracilis* have petal spots, as do other species in the section *Rhodanthos*, our analysis of spot formation in *C. gracilis* sheds light on aspects of how petal spots originally evolved in this group. Two major types of regulatory change appear to have been required for the evolution of spots. The first was a spatial restriction of the expression domain of *Dfr2*. It is reasonable to assume that pigment deposition throughout the petal was the ancestral state because species that are situated in the phylogeny as sister to the rest of the genus, namely *C. breweri*, *C. concinna*, and *C. pulchella*, are all pink and unspotted. Although it is unclear when the duplication of *Dfr* occurred, it is likely that initially both paralogs exhibited the same broad spatial expression pattern. Restriction of *Dfr2* expression to the areas of spot formation subsequently occurred. Because *Dfr2* seems no longer to be activated by the anthocyanin transcription factors that activate the other core anthocyanin enzymes, we suspect that its *cis*-regulatory region evolved to respond to a new set of activators that are expressed at the time and position at which spot formation initiates. The gene products of the *P* and *I* loci are candidates for these new activators, as they influence expression of *Dfr2* but not expression of the other core enzyme-coding genes.

The second major type of regulatory change necessary for the evolution of cyanidin-based spots separated the timing of expression of *Dfr2* and *F3'h* on the one hand from *Dfr1* and *F3'5'h* on the other. By itself, divergence in spatial expression domains of the two *Dfr* paralogs might produce a spot of increased pigment intensity, which could explain the existence of *Clarkia* species that produce spots but only malvidin in their flowers (Soltis, 1986), but cannot account for cyanidin/peonidin spots on a malvidin background, as seen in *C. gracilis*. Our analyses suggest that temporal separation of the expression of *F3'h* and *F3'5'h* is largely responsible for the production of cyanidin/peonidin-based anthocyanin in spots.



**Fig. 6** Model of pigment pattern development in *Clarkia gracilis* flowers. Early pigment production in spots is explained by flavonoid 3' hydroxylase (*F3'h*) expression throughout the petal and dihydroflavonol-4-reductase 2 (*Dfr2*) expression in spots, which leads to dihydroquercetin (DHQ) production throughout the petal and cyanidin/peonidin production in spots only. Later in development, expression of flavonoid 3'-5' hydroxylase (*F3'5'h*) and *Dfr1* converts DHQ (and any residual dihydrokaempferol (DHK)) in the petal background to dihydromyricetin (DHM) and then malvidin.

In most plant species that have been examined, both *F3'h* and *F3'5'h* are active simultaneously and result in the production of malvidin/delphinidin derivatives (Gerats *et al.*, 1982; Tornielli *et al.*, 2009; Hopkins & Rausher, 2011). We therefore presume that this was the pattern in ancestors of *C. gracilis* that lacked spots (only malvidin is produced in the unspotted flowers of *C. breweri* and *C. concinna*; Soltis, 1986). Cyanidin, peonidin and their derivatives can only be produced if *F3'5'h* activity is greatly reduced or eliminated. *Clarkia gracilis* has apparently achieved this by delaying the expression of *F3'5'h* so that, at the time of spot development, only *F3'h* is expressed. By contrast, at the time of background pigment development, only *F3'5'h* is expressed, allowing the accumulation of malvidin derivatives. Finally, restriction of the expression of *Dfr1* to the late stage of development was required to prevent cyanidin- and peonidin-based anthocyanins from being synthesized in the petal background.

Our analysis of the genetic control of spot pattern formation has revealed that the gene *Dfr2* acts as a switch for spot production. Activation of this gene leads to pigment deposition in spots. The evolution of this switch mechanism appears to have been facilitated by duplication of the ancestral *Dfr* gene, which allowed one paralog to evolve to serve as a spot regulator. Based on the broad occurrence of petal spots in *Clarkia*, we infer that the *Dfr* gene duplication occurred early in the radiation of the genus. Subsequent evolutionary remodeling of the regulatory network for both copies of *Dfr*, as well as for *F3'h* and *F3'5'h*, resulted in the novel ability to finely regulate the spatial pattern of contrasting colors in the petals of many *Clarkia* species. Further work in this group, particularly temporal and spatial expression assays of anthocyanin pathway genes in petals from species throughout the genus, will help to illuminate the order of events that led to the evolution of petal spot.

## Acknowledgements

We would like to thank the late Leslie Gottlieb for numerous helpful discussions and for providing us with his original seed materials. We thank two anonymous reviewers for their helpful suggestions on how to improve this paper. We also thank Brian Barringer and Norman Weeden for seeds, Ken Sytsma for DNA samples, and Vince Eckhart for primer sequences. Todd Barkman is thanked for helpful discussions. Kandis Elliot and Sara Friedrich are thanked for assistance with images. This work was supported by a NIH-Genetics Training Grant, a University of Wisconsin Sci-Med GRS Fellowship, a Ford Foundation Dissertation Fellowship, and a NSF DBI-1103693 award to TRM, and a University of Wisconsin Hamel Family Faculty Fellowship to DAB.

## References

- Abdel-Hameed F, Snow R. 1972. The origin of the allotetraploid *Clarkia gracilis*. *Evolution* 26: 74–83.
- Albert NW, Lewis DH, Zhang H, Schwinn KE, Jameson PE, Davies KM. 2011. Members of an R2R3-MYB transcription factor family in *Petunia* are developmentally and environmentally regulated to control complex floral and vegetative pigmentation patterning. *The Plant Journal* 65: 771–784.
- Beldade P, Brakefield PM, Long AD. 2002. Contribution of *Distal-less* to quantitative variation in butterfly eyespots. *Nature* 415: 315–318.
- Borevitz JO, Xia YJ, Blount J, Dixon RA, Lamb C. 2000. Activation tagging identifies a conserved MYB regulator of phenylpropanoid biosynthesis. *Plant Cell* 12: 2383–2393.
- Carey CC, Strahle JT, Selinger DA, Chandler VL. 2004. Mutations in the pale *aleurone color1* regulatory gene of the *Zea mays* anthocyanin pathway have distinct phenotypes relative to the functionally similar *TRANSPARENT TESTA GLABRA1* gene in *Arabidopsis thaliana*. *Plant Cell* 16: 450–464.
- Carroll SB, Gates J, Keys DN, Paddock SW, Panganiban GB, Selegue JE, Williams JA. 1994. Pattern formation and eyespot determination in butterfly wings. *Science* 265: 109–114.
- Cone K, Burr F, Burr B. 1986. Molecular analysis of the maize anthocyanin regulatory locus C1. *Proceedings of the National Academy of Sciences, USA* 83: 9631–9635.
- DesMarais D, Rausher MD. 2008. Escape from adaptive conflict after duplication in an anthocyanin pathway gene. *Nature* 454: 762–766.
- Dorn P, Bloom W. 1984. Anthocyanin variation in an introgressive complex in *Clarkia*. *Biochemical Systematics and Ecology* 12: 311–314.
- Doyle JJ, Doyle JL. 1987. A rapid DNA isolation procedure from small quantities of fresh leaf tissues. *Phytochemical Bulletin* 19: 11–15.
- Eckhart M, Rushing S, Hart M. 2006. Frequency dependent pollinator foraging in polymorphic *Clarkia xantiana* ssp. *xantiana* populations: implications for flower colour evolution and pollinator interactions. *Oikos* 112: 412–421.
- Elomaa P, Mehto M, Kotilainen M, Helariutta Y, Nevalainen L, Teeri T. 1998. A bHLH transcription factor mediates organ, region and flower type specific signals on dihydroflavonol-4-reductase (*dfr*) gene expression in the inflorescence of *Gerbera hybrida* (Asteraceae). *The Plant Journal* 16: 93–99.
- Fischer T, Halbwirth H, Meisel B, Stich K. 2003. Molecular cloning, substrate specificity of the functionally expressed dihydroflavonol 4-reductases from *Malus domestica* and *Pyrus communis* cultivars and the consequences for flavonoid metabolism. *Archives of Biochemistry and Biophysics* 412: 223–230.
- Gerats AGM, de Vlaming P, Doodeman M, Al B, Schram AW. 1982. Genetic control of the conversion of dihydroflavonols into flavonols and anthocyanins in flowers of *Petunia hybrida*. *Planta* 155: 364–368.
- Gonzalez A, Zhao M, Leavitt JM, Lloyd AM. 2008. Regulation of the anthocyanin biosynthetic pathway by the TTG1/bHLH/Myb transcriptional complex in *Arabidopsis* seedlings. *The Plant Journal* 53: 814–827.
- Goodrich J, Carpenter R, Coen ES. 1992. A common gene regulates pigmentation pattern in diverse plant species. *Cell* 68: 955–964.
- Gottlieb L, Ford V. 1988. Genetic studies of the pattern of floral pigmentation in *Clarkia gracilis*. *Heredity* 60: 237–246.
- Goulson D, Mcguire K, Munro EE, Adamson S, Collier L, Park KJ, Tinsley MC, Gilburn AS. 2009. Functional significance of the dark central floret of *Daucus carota* (Apiaceae) L.; is it an insect mimic? *Plant Species Biology* 24: 77–82.
- Harborne JB. 1984. *Phytochemical methods: a guide to modern techniques of plant analysis*, 2nd edn. London, UK: Chapman and Hall.
- Hopkins R, Rausher MD. 2011. Identification of two genes causing reinforcement in the Texas wildflower *Phlox drummondii*. *Nature* 469: 411–414.
- Jackson D, Roberts K, Martin C. 1992. Temporal and spatial control of expression of anthocyanin biosynthetic genes in developing flowers of *Antirrhinum majus*. *The Plant Journal* 2: 425–434.
- Johnson S, Midgley J. 1997. Fly pollination of *Gorteria diffusa* (Asteraceae), and a possible mimetic function for dark spots on the capitulum. *American Journal of Botany* 84: 429–436.
- Johnson E, Ryu S, Yi H, Shin B, Cheong H, Choi G. 2001. Alteration of a single amino acid changes the substrate specificity of dihydroflavonol-4-reductase. *The Plant Journal* 25: 325–333.
- Jones K. 1996. Pollinator behavior and postpollination reproductive success in alternative floral phenotypes of *Clarkia gracilis* (Onagraceae). *International Journal of Plant Sciences* 157: 733–738.
- Keys DN, Lewis DL, Selegue JE, Pearson BJ, Goodrich LV, Johnson RL, Gates J, Scott MP, Carroll SB. 1999. Recruitment of a hedgehog regulatory circuit in butterfly eyespot evolution. *Science* 283: 532–534.
- Lin-Wang K, Bolitho K, Grafton K, Kortstee A, Karunaratnam S, McGhie TK, Espley RV, Hellens RP, Allan AC. 2010. An R2R3 MYB transcription factor associated with regulation of the anthocyanin biosynthetic pathway in Rosaceae. *BMC Plant Biology* 10: 50.
- Ludwig SR, Habera LF, Dellaporta SL, Wessler SR. 1989. *Lc*, a member of the maize *R* gene family responsible for tissue-specific anthocyanin production,

- encodes a protein similar to transcriptional activators and contains the *Myc*-homology region. *Proceedings of the National Academy of Sciences, USA* 86: 7092–7096.
- Maddison DR, Maddison WP. 2000. *MacClade 4: analysis of phylogeny and character evolution*. Sunderland, MA, USA: Sinauer Associates.
- Martin C, Prescott A, Mackay S, Bartlett J, Vrijlandt E. 1991. Control of anthocyanin biosynthesis in flowers of *Antirrhinum majus*. *The Plant Journal* 1: 37–49.
- Morita Y, Saitoh M, Hoshino A, Nitasaka E, Iida S. 2006. Isolation of cDNAs for *R2R3-MYB*, *bHLH* and *WDR* transcriptional regulators and identification of *c* and *ca* mutations conferring white flowers in the Japanese morning glory. *Plant and Cell Physiology* 47: 457–470.
- Nijhout HF. 1980. Pattern formation on Lepidopteran wings: determination of an eyespot. *Developmental Biology* 80: 267–274.
- Niu S-S, Xu C-J, Zhang W-S, Zhang B, Li X, Lin-Wang K, Ferguson IB, Allan AC, Chen K-S. 2010. Coordinated regulation of anthocyanin biosynthesis in Chinese bayberry (*Myrica rubra*) fruit by a R2R3 MYB transcription factor. *Planta* 231: 887–899.
- Parker HG, VonHoldt BM, Quignon P, Margulies EH, Shao S, Mosher DS, Spady TC, Elkahoulou A, Cargill M, Jones PG *et al.* 2009. An expressed *Fgf4* retrogene is associated with breed-defining chondrodysplasia in domestic dogs. *Science* 325: 995–998.
- Paz-Ares J, Ghosal D, Wienand U, Peterson PA, Saedler H. 1987. The regulatory *c1* locus of *Zea mays* encodes a protein with homology to *myb* proto-oncogene products and with structural similarities to transcriptional activators. *EMBO Journal* 6: 3553–3558.
- Paz-Ares J, Wienand U, Peterson P, Saedler H. 1986. Molecular cloning of the *c* locus of *Zea mays*: a locus regulating the anthocyanin pathway. *EMBO Journal* 9: 315–321.
- Pichersky E, Raguso R, Lewinsohn E. 1994. Floral scent production in *Clarkia* (Onagraceae) (I. Localization and developmental modulation of monoterpene emission and linalool synthase activity). *Plant Physiology* 106: 1533–1540.
- Quattrocchio F, Wing J, van der Woude K, Mol J, Koes R. 1998. Analysis of bHLH and MYB domain proteins: species-specific regulatory differences are caused by divergent evolution of target anthocyanin genes. *The Plant journal* 13: 475–488.
- Quattrocchio F, Wing J, van der Woude K, Souer E, de Vetten N, Mol J, Koes R. 1999. Molecular analysis of the *anthocyanin2* gene of petunia and its role in the evolution of flower color. *Plant Cell* 11: 1433–1444.
- Reed RD, Serfas MS. 2004. Butterfly wing pattern evolution is associated with changes in a Notch/Distal-less temporal pattern formation process. *Current Biology* 14: 1159–1166.
- Robinson G, Robinson R. 1931. CLXXXII. A survey of anthocyanins. I. *Biochemical Journal* 25: 1687–1705.
- Schwinn K, Venail J, Shang Y, Mackay S, Alm V. 2006. A small family of MYB-regulatory genes controls floral pigmentation intensity and patterning in the genus *Antirrhinum*. *The Plant Cell* 18: 831–851.
- Shang Y, Venail J, Mackay S, Bailey PC, Schwinn KE, Jameson PE, Martin CR, Davies KM. 2011. The molecular basis for venation patterning of pigmentation and its effect on pollinator attraction in flowers of *Antirrhinum*. *New Phytologist* 189: 602–615.
- Shimada N, Sasaki R, Sato S. 2005. A comprehensive analysis of six dihydroflavonol 4-reductases encoded by a gene cluster of the *Lotus japonicus* genome. *Journal of Experimental Botany* 56: 2573–2585.
- Smith S, Rauscher MD. 2011. Gene loss and parallel evolution contribute to species difference in flower color. *Molecular Biology and Evolution* 28: 2799–2810.
- Soltis P. 1986. Anthocyanidin variation in *Clarkia*. *Biochemical Systematics and Ecology* 14: 487–489.
- Spelt C, Quattrocchio F, Mol JNM, Koes R. 2000. *Anthocyanin1* of petunia encodes a basic helix-loop-helix protein that directly activates transcription of structural anthocyanin genes. *Plant Cell* 12: 1619–1631.
- Spelt C, Quattrocchio F, Mol J, Koes R. 2002. ANTHOCYANIN1 of petunia controls pigment synthesis, vacuolar pH, and seed coat development by genetically distinct mechanisms. *Plant Cell* 14: 2121–2135.
- Swofford DL. 2002. *PAUP\*: phylogenetic analysis using parsimony (\* and other methods), version 4.0b10*. Sunderland, MA, USA: Sinauer Associates.
- Tornielli G, Koes R, Quattrocchio F. 2009. The genetics of flower color. In: Gerats T, Strommer J, eds. *Petunia, evolutionary, developmental and physiological genetics*. New York, NY, USA: Springer, 269–299.
- Van Kleunen M, Nänni I, Donaldson J, Manning J. 2007. The role of beetle marks and flower colour on visitation by monkey beetles (Hopliini) in the greater cape floral region, South Africa. *Annals of Botany* 100: 1483–1489.
- de Vetten N, Quattrocchio F, Mol J, Koes R. 1997. The *an11* locus controlling flower pigmentation in petunia encodes a novel WD-repeat protein conserved in yeast, plants, and animals. *Genes & Development* 11: 1422–1434.
- Walker AR. 1999. The *TRANSPARENT TESTA GLABRA1* Locus, which regulates trichome differentiation and anthocyanin biosynthesis in *Arabidopsis*, encodes a WD40 Repeat protein. *Plant Cell* 11: 1337–1349.
- Xiao H, Jiang N, Schaffner E, Stockinger EJ, van der Knaap E. 2008. A retrotransposon-mediated gene duplication underlies morphological variation of tomato fruit. *Science* 319: 1527–1530.

## Supporting Information

Additional supporting information may be found in the online version of this article.

**Fig. S1** Phenotypes and sections of petals used for HPLC analyses.

**Fig. S2** Phenotypes, ages, and sections of petals used for qualitative RT-PCR analyses.

**Fig. S3** Stages of *Clarkia gracilis* petal development.

**Fig. S4** Genotypes at the *P* and *I* loci and expected phenotypes.

**Fig. S5** Phylogenetic trees of anthocyanin pathway genes in section *Rhodanthos*.

**Fig. S6** Dihydroflavonol-4-reductase 2 *A/B* (*Dfr2-A/B*) expression and *Actin* control in *C. gracilis* plants derived from seeds collected in natural populations.

**Fig. S7** Alignment of the six different dihydroflavonol-4-reductase (*Dfr*) sequences found in *Clarkia gracilis*.

**Table S1** Voucher information

**Table S2** Primers used in qualitative and quantitative PCR assays

**Table S3** Primers used and expected sizes for dihydroflavonol-4-reductase 2 (*Dfr2*) genotype assays for the *P* cross

**Table S4** Primers used and expected sizes for dihydroflavonol-4-reductase 2 (*Dfr2*) genotype assays for the *I* cross

**Notes S1** Text files containing alignments used for phylogenetic analyses.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.